

Meadowscape Pro™

Native Enriched

Product Specification

Contents

Technical Specification	2
SpeciesList.....	3&4

Appendix

Installation Guide	5
Maintenance Plan	6

Product Code:

MDSCNAT

Manufacturer:

Wildflower Co.

Manufacturer Contact Details

☎ 01256 771 222

✉ wildflower@wildflowerexperts.com

🌐 www.wildflowerexperts.com

Technical Specification

Product code	MDSCNAT
Manufacturer	Wildflower Co.
Dimensions	Delivered in 1m ³ bulk bags or 2m ³ bulk bags equating to 40m ² or 80m ²
Weight	400kg for a 1m ³ bulk bag and 800kg for a 2m ³ bulk bag
No. of species	54 species (UK provenance native and naturalised species) Access full species list here.
Grass:flora sowing ratio	100% wildflowers. Includes some annuals.
Flowering period	March-October
Established height	30-150 cm (Avg height 70cm).
Time to establish	3-12 months (at least 1 year for plants that need vernalisation period)
Soil type	Suitable for all soil types
Seed provenance and testing	Grown in England using UK native and naturalised species. Seed analysis results available on request.
Recyclability	100%, including packaging.
BNG suitability	Not suitable to BNG projects
Substrate features	Contains Hydropor™ for greater moisture retention
Seed mix	Calibrated using seed weight analysis

Scientifically Engineered: About this pre-seeded growing medium

This revolutionary substrate isn't your average soil. It's been scientifically engineered and rigorously tested to be the perfect incubator for the wildflower seeds. Compared to traditional methods, this translates to a dramatically improved germination process.

Our unique, 100% natural and peat-free formula slashes the full establishment time from years for traditional seeding, to just 3-12 months, depending on sowing season.

Our seed mixes are already integrated. They are meticulously formulated based on scientific analysis. Seed weight, size and germination undergoes lab analysis for optimum inclusion rates. Omitting this process affects species diversity.

* Actual % may vary, based on environmental factors and seed availability
You may not see every species listed as the product is designed to adapt to individual settings, including a wide range of soil types and environmental conditions. Many species may not flower within the first year.

Meadscape Pro™ Native Enriched

Product code: MDSCNAT

Get a Quote: www.wildflowerexperts.com

Species List

Meadscape Pro™ Native Enriched

Product code: MDSCNAT

All seed grown by Wildflower Co. is tested to stringent standards for germination and purity.

Seed analysis results are available on request.

*You may not see every species listed as the product is designed to adapt to individual settings, including a wide range of soil types and environmental conditions. Many species may not flower during the first year.

Species	Flowering Time	Ultimate Height	Colour	Life Cycle
Wildflower Species				
Autumn Hawkbit (<i>Scorzoneroides autumnalis</i>)	June - Oct	50	Yellow	P
Babys Breath (<i>Gypsophila elegans</i>)	June - Aug	100	White	A
Betony (<i>Betonica officinalis</i>)	June - Aug	60	Pink	P
Birdsfoot Trefoil (<i>Lotus corniculatus</i>)	May - Sept	30	Yellow	P
Bladder Campion (<i>Silene vulgaris</i>)	May - Sept	100	White	P
Borage (<i>Borago officinalis</i>)	May - Sept	60	Purple	A
Cats Ear (<i>Hypochaeris radicata</i>)	June - Sept	45	Yellow	P
Common/Cottage Pink (<i>Dianthus plumarius</i>)	May - July	50	Pink	P
Common Knapweed (<i>Centaurea nigra</i>)	June - Sept	100	Purple	P
Common Toadflax (<i>Linaria vulgaris</i>)	June - Nov	75	Yellow	P
Common Vetch (<i>Vicia sativa</i>)	Apr - Sept	75	Purple	A
Common Poppy (<i>Papaver rhoeas</i>)	June - Aug	60	Red	A
Corn Chamomile (<i>Anthemis arvensis</i>)	June - Sept	30	White	A
Corn Marigold (<i>Glebionis segetum</i>)	June - Oct	100	Yellow	A
Corncockle (<i>Agrostemma githago</i>)	May - Sept	120	Pink	A
Cornflower (<i>Centaurea cyanus</i>)	June - Sept	100	Blue	A
Cowslip (<i>Primula veris</i>)	Apr - June	25	Yellow	P
Crimson Clover (<i>Trifolium incarnatum</i>)	June - Sept	50	Red	A
Dames-violet (<i>Hepseris matronalis</i>)	May - July	90	Purple	BI
English (Pot) Marigold (<i>Calendula officinalis</i>)	June - Oct	50	Orange	BI
Field Forget-me-not (<i>Myosotis arvensis</i>)	Apr - Sept	40	Blue	SLP
Field Scabious (<i>Knautia arvensis</i>)	July - Oct	150	Purple	P
Golden Candytuft (<i>Iberis sempervirens</i>)	Apr - May	30	White	A
Great/Large Bellflower (<i>Campanula latifolia</i>)	June - Aug	120	Purple	P
Greater Knapweed (<i>Centaurea scabiosa</i>)	June - Sept	100	Purple	P
Hedge Bedstraw (<i>Galium album</i>) (prev. <i>mollugo</i>)	June - Sept	100	White	P
Hoary Plantain (<i>Plantago media</i>)	May - Aug	30	White	P
Kidney Vetch (<i>Anthyllis vulneraria</i>)	June - Sept	25	Yellow	P
Lady's Bedstraw (<i>Galium verum</i>)	June - Sept	50	Yellow	P
Larkspur (<i>Delphinium consolida</i>)	June - Sept	120	Pink	A
Meadow Cranesbill (<i>Geranium pratense</i>)	June - Sept	90	Purple	P
Mountain Madwort (<i>Alyssum montanum</i>)	May - July	15	Yellow	P
Musk Mallow (<i>Malva moschata</i>)	July - Aug	100	Purple	P
Nodding Catchfly (<i>Silene pendula</i>)	June - Sept	50	Pink	A
Oxeye Daisy (<i>Leucanthemum vulgare</i>)	May - July	90	White	P
Perforate St John's Wort (<i>Hypericum perforatum</i>)	June - Aug	150	Yellow	P
Perrenial Flax (<i>Linum perenne</i>)	May - June	60	Purple	P
Ragged Robin (<i>Silene flos-cuculi</i>)	June - Aug	100	Pink	P
Red Campion (<i>Silene dioica</i>)	Apr - July	100	Pink	P
Ribwort Plantain (<i>Plantago lanceolata</i>)	Apr - Oct	90	White	P
Sainfoin (<i>Onobrychis vicifolia</i>)	June - Aug	80	Pink	P
Salad Burnet (<i>Sanguisorba minor</i>)	June - Aug	65	Red	P
Self Heal (<i>Prunella vulgaris</i>)	May - Sept	30	Purple	P

Get a Quote: www.wildflowerexperts.com

Species List

Meadscape Pro™ Native Enriched

Product code: MDSCNAT

All seed grown by Wildflower Co. is tested to stringent standards for germination and purity.

Seed analysis results are available on request.

Species	Flowering Time	Ultimate Height	Colour	Life Cycle
Wildflower Species				
Sweet William (<i>Dianthus barbatus</i>)	June - Sept	80		BI / SLP
Sweet William Catchfly (<i>Silene armeria</i>)	May - Sept	45		P
Tufted Vetch (<i>Vicia cracca</i>)	June - Aug	120		P
Viper's Bugloss (<i>Echium vulgare</i>)	June - Aug	120		BI
Wallflower (<i>Erysimum cheiri</i>)	Mar - June	50		BI / P
White Campion (<i>Silene latifolia</i>)	May - Sept	100		SLP
Wild Marjoram (<i>Origanum vulgare</i>)	June - Aug	100		P
Wild Red Clover (<i>Trifolium pratense</i>)	May - Sept	50		P
Woodland Sage (<i>Salvia nemorosa</i>)	June - Sept	80		P
Yarrow (<i>Achillea millefolium</i>)	June - Aug	90		P
Yellow Rattle (<i>Rhinanthus minor</i>)	May - July	50		SLP

*You may not see every species listed as the product is designed to adapt to individual settings, including a wide range of soil types and environmental conditions. Many species may not flower during the first year.

Get a Quote: www.wildflowerexperts.com

Preparation and installation

Meadowscape Pro™

Soil preparation

Do NOT rotovate, dig or disturb the soil. The existing vegetation should be killed or removed. If spraying off the area, this should be conducted according to the chemical manufacturer's instructions a minimum of 2 weeks before installation of the Meadowscape Pro™. If there is a residue of more than a light covering of dead surface vegetation after two weeks then this needs to be cut and removed. This is necessary otherwise Meadowscape Pro™ will be suspended on the dead thatch above the soil surface, preventing good seed germination and successful seedling growth. Meadowscape Pro™ can also be laid on sterile imported soil, but if so the soil needs to be levelled for a smooth finish and for ease of laying product.

Soil conditions and fertility

Ensure soil is not waterlogged or compacted prior to installing Meadowscape Pro™. The soil does not need to be fertilised before or after the installation. We would advise not stripping back the top soil to reduce soil fertility before using Meadowscape Pro™ as it needs some level of fertility to get well-established initially and is an unnecessary ground preparation step. However, where soil is fertile, particular attention must be paid to the maintenance regime – see maintenance section.

Laying Meadowscape Pro™

When laying Meadowscape Pro™, care should be taken to ensure a uniform depth of 20 – 25mm is laid. Do not over or under apply the material. Over application results in an imbalance of seed and nutrient ratios for a given area creating too much competition between the plants. Under applying will create a thin layer of seeded material creating sparser and poorer germination results whilst also limiting Meadowscape Pro™'s ability to suppress weed seeds in the surface of the soil beneath.

Watering

Once laid, water the area thoroughly, for the first few weeks until a seedling sward starts to appear– this could take up to 4-6 weeks (weather dependent). There is no need to soak the soil before laying but it is important to check that the initial watering soaks through the entire dressing of growing medium and ideally into the soil beneath. Do this by scratching the growing medium during watering to assess sufficient water uptake.

Once the Meadowscape Pro™ is wet through, stop watering – excessive watering will lead to run off and erosion of the material. Do not allow the profile of the Meadowscape Pro™ to dry out while the seeds establish, it need to be damp – this is very important. Depending on the time of year, seedlings will emerge in the first few weeks and these plants will develop rapidly as a result of the perfect growing conditions the Meadowscape Pro™ provides.

Meadowscape Pro™ Native Enriched

Product code: MDSCNAT

Get a Quote: www.wildflowerexperts.com

Maintenance

Management plan for long term success

Fertiliser

No fertiliser is needed, although in some circumstances, for example where the wildflower installation is on very low fertility soil such as sand or gravel, the addition of a light dose of fertiliser in the spring may improve plant development. Please contact Wildflower Co. for guidance.

Annual Maintenance

Once established Meadowscape Pro™ requires little maintenance. For the annual maintenance cut in the autumn, it is important to cut the meadow down to 2-3 inches (50-75mm) off the ground and remove all cuttings. This can be done by strimming and raking, or using a mower and collecting the cuttings. Make sure these tools are sharp. The cut is an important part of the meadows life cycle and ensures that re-growth and species diversity improves year on year. Cuttings should not be left on the meadow, as they add undesirable fertility to the ground. It is also important to remove all leaf litter that falls onto the area.

Timing the cut

The annual maintenance cut should be done in late September, early October. There is no need for a set date, but this timing will allow the plants in the meadow to regenerate before the first frost typically in November. You can choose to cut only half of the meadow area at one time to allow time for fauna to migrate to the uncut meadow. Allow some regrowth of the cut area before cutting the second half, but aim to have finished all cutting by the end of the first week of October. Over time alternate the areas that are cut early and the areas that are left as this will benefit species diversity.

Managing the fertility in the ground

On fertile sites or where you might have species dominance or too vigorous early growth, a second cut at the end of May, beginning of June can be introduced. This high cut, approximately 8 to 10 inches off the ground (200mm to 250mm) to remove the flower heads but leaving enough plant stems and leaf area to regenerate, and removal of all cuttings, will help to knock back some species dominance, reduce soil fertility and open up the sward to more light and air circulation to promote diversity of lower growing species. Once the cutting has been completed and all cuttings removed, give the area a good soaking with water to encourage the next flush of growth. Introducing this early summer cut and removal will mean your second autumn cut and removal will be later that year, up to the end of October.

Meadowscape Pro™ Native Enriched

Product code: MDSCNAT